
A review of classical propositional logic



Chrysippus, c. 279 BC - c. 206 BC.

We start with a (finite or infinite) set of propositional variables, or atomic formulæ, that are to stand for propositions. Say, if we content ourselves with countably many:

$$X_1, X_2, \ldots, X_n, \ldots$$

(We can use p, q, etc. as a lighter short-hand notation.)

We start with a (finite or infinite) set of propositional variables, or atomic formulæ, that are to stand for propositions. Say, if we content ourselves with countably many:

$$X_1, X_2, \ldots, X_n, \ldots$$

(We can use p, q, etc. as a lighter short-hand notation.) To these we adjoin two symbols \top and \bot , say, that are to stand for a proposition that is always true (the *verum*), and one that is always false (the *falsum*), respectively.

We start with a (finite or infinite) set of propositional variables, or atomic formulæ, that are to stand for propositions. Say, if we content ourselves with countably many:

$$X_1, X_2, \ldots, X_n, \ldots$$

(We can use p, q, etc. as a lighter short-hand notation.) To these we adjoin two symbols \top and \bot , say, that are to stand for a proposition that is always true (the verum), and one that is always false (the falsum), respectively.

To construct compound formulæ we use the logical connectives:

- ∨, for disjunction ("inclusive or", Latin vel, unlike Latin aut);
- ∧, for conjunction ("and", Latin et);
- \blacksquare \rightarrow , for implication ("if...then...", conditional assertions);
- ¬, for negation ("not", negative assertions).

A review of classical propositional logic

A review of classical propositional logic

The usual recursive definition of general formulæ now reads as follows.

 \blacksquare \top and \bot are formulæ.

- \blacksquare \top and \bot are formulæ.
- All propositional variables are formulæ.

- \blacksquare \top and \bot are formulæ.
- All propositional variables are formulæ.
- If α and β are formulæ, so are $(\alpha \lor \beta)$, $(\alpha \land \beta)$, $(\alpha \to \beta)$, and $\neg \alpha$.

- \blacksquare \top and \bot are formulæ.
- All propositional variables are formulæ.
- If α and β are formulæ, so are $(\alpha \lor \beta)$, $(\alpha \land \beta)$, $(\alpha \to \beta)$, and $\neg \alpha$.
- Nothing else is a formula.

- \blacksquare \top and \bot are formulæ.
- All propositional variables are formulæ.
- If α and β are formulæ, so are $(\alpha \lor \beta)$, $(\alpha \land \beta)$, $(\alpha \to \beta)$, and $\neg \alpha$.
- Nothing else is a formula.

Let us write FORM for the set of all formulæ constructed over the countable language X_1, \ldots, X_n, \ldots , and FORM_n for the set of formulæ constructed over the first n propositional variables X_1, \ldots, X_n , at the most.

A review of classical propositional logic

We now construct a formal semantics in order to interpret formulæ.

A review of classical propositional logic

We now construct a formal semantics in order to interpret formulæ. Namely, we consider assignments of truth-values, or evaluations, or interpretations.

These are functions

$$w \colon \text{Form} \to \{0, 1\}$$

These are functions

$$w \colon \text{Form} \to \{0, 1\}$$

$$w(\top) = 1, w(\bot) = 0;$$

These are functions

$$w \colon \text{Form} \to \{0, 1\}$$

- $w(\top) = 1, w(\bot) = 0;$
- $w(\alpha \land \beta) = 1$ if and only if both $w(\alpha) = 1$ and $w(\beta) = 1$;

These are functions

$$w \colon \text{Form} \to \{0, 1\}$$

- $w(\top) = 1, w(\bot) = 0;$
- $w(\alpha \land \beta) = 1$ if and only if both $w(\alpha) = 1$ and $w(\beta) = 1$;
- $w(\alpha \lor \beta) = 1$ if and only if either $w(\alpha) = 1$ or $w(\beta) = 1$ (or both);

These are functions

$$w: \text{Form} \rightarrow \{0, 1\}$$

- $w(\top) = 1, w(\bot) = 0;$
- $w(\alpha \wedge \beta) = 1$ if and only if both $w(\alpha) = 1$ and $w(\beta) = 1$;
- $w(\alpha \lor \beta) = 1$ if and only if either $w(\alpha) = 1$ or $w(\beta) = 1$ (or both);
- $w(\alpha \to \beta) = 0$ if and only if $w(\alpha) = 1$ and $w(\beta) = 0$;

These are functions

$$w \colon \text{Form} \to \{0, 1\}$$

- $w(\top) = 1, w(\bot) = 0;$
- $w(\alpha \land \beta) = 1$ if and only if both $w(\alpha) = 1$ and $w(\beta) = 1$;
- $w(\alpha \lor \beta) = 1$ if and only if either $w(\alpha) = 1$ or $w(\beta) = 1$ (or both);
- $w(\alpha \to \beta) = 0$ if and only if $w(\alpha) = 1$ and $w(\beta) = 0$;
- $w(\neg \alpha) = 1$ if and only if $w(\alpha) = 0$.

A review of classical propositional logic

A first easy observation about these definitions is that a valuation w is subject to no restrictions concerning the values it assigns to propositional variables.

Moreover, it is also apparent that any valuation w is uniquely determined by such values $w(X_1), \ldots, w(X_n), \ldots$

Moreover, it is also apparent that any valuation w is uniquely determined by such values $w(X_1), \ldots, w(X_n), \ldots$

This is the principle of truth-functionality, or compositionality (after G. Frege) for classical propositional logic.

Moreover, it is also apparent that any valuation w is uniquely determined by such values $w(X_1), \ldots, w(X_n), \ldots$

This is the principle of truth-functionality, or compositionality (after G. Frege) for classical propositional logic.

Following a well-established tradition going back at least to G. W. Leibniz, it is useful to think of an evaluation $w \colon \mathrm{FORM} \to \{0,1\}$ as a possible world in which each proposition α either holds — i.e., $w(\alpha) = 1$ — or fails to hold – i.e., $w(\alpha) = 0$.

Moreover, it is also apparent that any valuation w is uniquely determined by such values $w(X_1), \ldots, w(X_n), \ldots$

This is the principle of truth-functionality, or compositionality (after G. Frege) for classical propositional logic.

Following a well-established tradition going back at least to G. W. Leibniz, it is useful to think of an evaluation $w \colon \mathrm{Form} \to \{0,1\}$ as a possible world in which each proposition α either holds — i.e., $w(\alpha) = 1$ — or fails to hold – i.e., $w(\alpha) = 0$.

Because of truth-functionality, then, a classical logician's world is something very ethereal indeed: it is uniquely determined by the collection of true atomic propositions that can be uttered.

$$lacksquare oxedsymbol{\perp}
ightarrow lpha$$
 (Ex falso quodlibet)

$$lacktriangledown$$
 $\alpha \lor \neg \alpha$ (Tertium non datur)

$$\qquad \qquad -(\alpha \wedge \neg \alpha) \qquad \qquad \text{(Principle of non-contradiction)}$$

| $lacksquare$ $oxdot$ \rightarrow $lpha$ | (Ex falso quodlibet) |
|--|----------------------------------|
| $\alpha \lor \neg \alpha$ | (Tertium non datur) |
| $\blacksquare \neg (\alpha \land \neg \alpha)$ | (Principle of non-contradiction) |
| lacksquare | (Law of double negation) |

$$\perp \rightarrow \alpha$$

$$\alpha \vee \neg \alpha$$

$$\neg (\alpha \land \neg \alpha)$$

$$\neg \neg \alpha \rightarrow \alpha$$

$$(\neg \alpha \rightarrow \alpha) \rightarrow \alpha$$

(Consequentia mirabilis)

$$\perp \rightarrow \alpha$$

$$\alpha \vee \neg \alpha$$

$$\neg (\alpha \land \neg \alpha)$$

$$\neg \neg \alpha \rightarrow \alpha$$

$$(\neg \alpha \rightarrow \alpha) \rightarrow \alpha$$

(Ex falso quodlibet)

(Tertium non datur)

(Principle of non-contradiction)

(Law of double negation)

(Consequentia mirabilis)

(Contraposition)

$$\perp \rightarrow \alpha$$

$$\alpha \vee \neg \alpha$$

$$\blacksquare \neg (\alpha \land \neg \alpha)$$

$$\neg \neg \alpha \rightarrow \alpha$$

$$(\neg \alpha \rightarrow \alpha) \rightarrow \alpha$$

Define: TAUT \subseteq FORM is the set of all tautologies. Similarly for TAUT_n \subseteq FORM_n.

Define: TAUT \subseteq FORM is the set of all tautologies. Similarly for TAUT_n \subseteq FORM_n.

Write: $\models \alpha$ to mean $\alpha \in TAUT$.

A review of classical propositional logic

Tautologies are a (formal) semantical notion. Logic is concerned with the relationship between syntax (the language) and semantics (the world).

Tautologies are a (formal) semantical notion. Logic is concerned with the relationship between syntax (the language) and semantics (the world).

The syntactic counterpart of a tautology is a provable formula, also called theorem of the logic.

Tautologies are a (formal) semantical notion. Logic is concerned with the relationship between syntax (the language) and semantics (the world).

The syntactic counterpart of a tautology is a provable formula, also called theorem of the logic.

To define provability, we select (with a lot of hindsight) a set of tautologies, and declare that they are axioms: they count as provable formulæ by definition.

Tautologies are a (formal) semantical notion. Logic is concerned with the relationship between syntax (the language) and semantics (the world).

The syntactic counterpart of a tautology is a provable formula, also called theorem of the logic.

To define provability, we select (with a lot of hindsight) a set of tautologies, and declare that they are axioms: they count as provable formulæ by definition.

Next we select a set of deduction rules that tell us that if we already established that formulæ $\alpha_1, \ldots, \alpha_n$ are provable, and these have a certain shape, then a specific formula β is also a provable formula.

$$rac{lpha \qquad lpha
ightarrow eta}{eta}$$
 (MP)

$$\frac{\alpha \qquad \alpha \to \beta}{\beta} \quad \text{(MP)}$$

Now we declare that a formula $\alpha \in \mathrm{FORM}$ is provable if there exists a proof of α , that is, a finite sequence of formulæ $\alpha_1, \ldots, \alpha_l$ a such that:

$$\frac{\alpha \qquad \alpha \to \beta}{\beta} \quad \text{(MP)}$$

Now we declare that a formula $\alpha \in \mathrm{FORM}$ is provable if there exists a proof of α , that is, a finite sequence of formulæ $\alpha_1, \ldots, \alpha_l$ a such that:

$$\alpha_l = \alpha$$
.

$$\frac{\alpha \qquad \alpha \to \beta}{\beta} \quad \text{(MP)}$$

Now we declare that a formula $\alpha \in \text{FORM}$ is provable if there exists a proof of α , that is, a finite sequence of formulæ $\alpha_1, \ldots, \alpha_l$ a such that:

- $\alpha_l = \alpha$.
- Each α_i , i < l is either an axiom, or is obtainable from α_j and α_k , j, k < i, via an application of *modus ponens*.

$$\frac{\alpha \qquad \alpha \to \beta}{\beta}$$
 (MP)

Now we declare that a formula $\alpha \in \text{FORM}$ is provable if there exists a proof of α , that is, a finite sequence of formulæ $\alpha_1, \ldots, \alpha_l$ a such that:

- $\alpha_l = \alpha$.
- Each α_i , i < l is either an axiom, or is obtainable from α_j and α_k , j, k < i, via an application of *modus ponens*.

Write: $\vdash \alpha$ if α is provable. Write: $\mathrm{THM} \subseteq \mathrm{FORM}$ for the set of provable formulæ. Similarly for $\mathrm{THM}_n \subseteq \mathrm{FORM}_n$.

$$\frac{\alpha \qquad \alpha \to \beta}{\beta} \quad \text{(MP)}$$

Now we declare that a formula $\alpha \in \text{FORM}$ is provable if there exists a proof of α , that is, a finite sequence of formulæ $\alpha_1, \ldots, \alpha_l$ a such that:

- $\alpha_l = \alpha$.
- Each α_i , i < l is either an axiom, or is obtainable from α_j and α_k , j, k < i, via an application of *modus ponens*.

Write: $\vdash \alpha$ if α is provable. Write: $\mathrm{THM} \subseteq \mathrm{FORM}$ for the set of provable formulæ. Similarly for $\mathrm{THM}_n \subseteq \mathrm{FORM}_n$.

We still need to define the axioms.

(A0)
$$\perp \rightarrow \alpha$$
 Ex falso quodlibet.

(A0)
$$\perp \rightarrow \alpha$$

(A1) $\alpha \rightarrow (\beta \rightarrow \alpha)$

Ex falso quodlibet.

A fortiori.

(A0)
$$\perp \rightarrow \alpha$$
 Ex falso quodlibet.
(A1) $\alpha \rightarrow (\beta \rightarrow \alpha)$ A fortiori.

(A2)
$$(\alpha \to \beta) \to ((\beta \to \gamma) \to (\alpha \to \gamma))$$
 Implication is transitive.

(A0)
$$\perp \rightarrow \alpha$$
 Ex falso quodlibet.
(A1) $\alpha \rightarrow (\beta \rightarrow \alpha)$ A fortiori.

(A2)
$$(\alpha \to \beta) \to ((\beta \to \gamma) \to (\alpha \to \gamma))$$
 Implication is transitive.

(A3)
$$((\alpha \rightarrow \beta) \rightarrow \beta) \rightarrow ((\beta \rightarrow \alpha) \rightarrow \alpha)$$

(A4)
$$(\alpha \to \beta) \to (\neg \beta \to \neg \alpha)$$
 Contraposition.

(A0)
$$\perp \rightarrow \alpha$$
 Ex falso quodlibet.

(A1)
$$\alpha \rightarrow (\beta \rightarrow \alpha)$$
 A fortiori.

(A2)
$$(\alpha \to \beta) \to ((\beta \to \gamma) \to (\alpha \to \gamma))$$
 Implication is transitive.

(A3)
$$((\alpha \rightarrow \beta) \rightarrow \beta) \rightarrow ((\beta \rightarrow \alpha) \rightarrow \alpha)$$

(A4)
$$(\alpha \to \beta) \to (\neg \beta \to \neg \alpha)$$
 Contraposition.

(A5)
$$(\neg \alpha \to \alpha) \to \alpha$$
 Consequentia Mirabilis.

Upon defining

$$\alpha \vee \beta \equiv (\alpha \rightarrow \beta) \rightarrow \beta$$

(A0-A5) read as shown next.

$$\alpha \vee \beta \equiv (\alpha \rightarrow \beta) \rightarrow \beta$$

(A0)
$$\perp \rightarrow \alpha$$

(A1)
$$\alpha \rightarrow (\beta \rightarrow \alpha)$$

A fortiori.

(A2)
$$(\alpha \to \beta) \to ((\beta \to \gamma) \to (\alpha \to \gamma))$$
 Implication is transitive.

(A3)
$$(\alpha \vee \beta) \rightarrow (\beta \vee \alpha)$$

Disjunction is commutative.

(A4)
$$(\alpha \to \beta) \to (\neg \beta \to \neg \alpha)$$

Contraposition.

(A5)
$$\alpha \vee \neg \alpha$$

Tertium non datur.

$$\alpha \vee \beta \equiv (\alpha \rightarrow \beta) \rightarrow \beta$$

Deduction rule for classical logic.

(R1)
$$\frac{\alpha \qquad \alpha \rightarrow \beta}{\beta}$$

Modus ponens.

This axiomatisation only uses \bot , \to , and \neg as primitive connectives. The remaining ones are definable as:

This axiomatisation only uses \bot , \rightarrow , and \neg as primitive connectives. The remaining ones are definable as:

$$\bullet \ \alpha \wedge \beta \ \equiv \ \neg (\neg \alpha \vee \neg \beta)$$
 (De Morgan's law)

$$\blacksquare \ \alpha \vee \beta \ \equiv \ (\alpha \to \beta) \ \to \ \beta$$

$$\bullet \ \alpha \wedge \beta \ \equiv \ \neg (\neg \alpha \vee \neg \beta)$$
 (De Morgan's law)

More defined connectives:

$$\bullet \ \alpha \wedge \beta \ \equiv \ \neg (\neg \alpha \vee \neg \beta)$$
 (De Morgan's law)

More defined connectives:

$$\bullet \ \alpha \leftrightarrow \beta \ \equiv \ (\alpha \to \beta) \land (\beta \to \alpha) \tag{Biconditional}$$

$$\bullet \ \alpha \wedge \beta \ \equiv \ \neg (\neg \alpha \vee \neg \beta)$$
 (De Morgan's law)

More defined connectives:

$$\bullet \ \alpha \leftrightarrow \beta \ \equiv \ (\alpha \to \beta) \land (\beta \to \alpha) \tag{Biconditional}$$

$$\bullet \ \alpha \ominus \beta \ \equiv \ \neg(\alpha \to \beta) \qquad \qquad \text{(Difference, or Co-implication)}$$

$$\bullet \ \alpha \wedge \beta \ \equiv \ \neg (\neg \alpha \vee \neg \beta)$$
 (De Morgan's law)

More defined connectives:

$$lacktriangledown$$
 $lpha \ominus eta \equiv \neg(lpha
ightarrow eta)$ (Difference, or Co-implication)

Exercise. Show that $\vdash \neg \neg \alpha \to \alpha$. Then pause to appreciate why proof-theorists develop Gentzen-style calculi to replace Hilbert-style calculi such as the one above.

Easy to show:

Easy to show:

Theorem (Validity or Soundness Theorem for CL)

For any $\alpha \in \mathrm{FORM}$,

 $\vdash \alpha \text{ implies } \models \alpha.$

That is,

Thm \subseteq Taut.

(Similarly, $THM_n \subseteq TAUT_n$).

Harder to show:

Harder to show:

Theorem (Completeness Theorem for CL)

For any $\alpha \in \mathrm{FORM}$,

$$\models \alpha \text{ implies } \vdash \alpha$$
.

That is,

Taut
$$\subseteq$$
 Thm.

(Similarly, $TAUT_n \subseteq THM_n$).

In summary:

In summary:

Theorem (Soundness and Completeness Theorem for CL)

For any $\alpha \in FORM$,

$$\models \alpha$$
 if, and only if, $\vdash \alpha$.

That is,

$$TAUT = THM$$
.

(Similarly, $TAUT_n = THM_n$).

Immediate consequence:

Immediate consequence:

Corollary (Decidability of CL)

There is a an algorithm (for definiteness, a Turing machine) that, on input $\alpha \in \mathrm{FORM}$, outputs YES if $\alpha \in \mathrm{THM}$, and NO otherwise.

Immediate consequence:

Corollary (Decidability of CL)

There is a an algorithm (for definiteness, a Turing machine) that, on input $\alpha \in \mathrm{FORM}$, outputs YES if $\alpha \in \mathrm{THM}$, and NO otherwise.

In light of completeness, one such algorithm is: check all possible $\{0,1\}$ -valued assignments to the n variables occurring in α . Since there is 2^n of them, computationally this is a pretty silly algorithm.

Immediate consequence:

Corollary (Decidability of CL)

There is a an algorithm (for definiteness, a Turing machine) that, on input $\alpha \in \mathrm{FORM}$, outputs YES if $\alpha \in \mathrm{THM}$, and NO otherwise.

In light of completeness, one such algorithm is: check all possible $\{0,1\}$ -valued assignments to the n variables occurring in α . Since there is 2^n of them, computationally this is a pretty silly algorithm.

An yet, it is hard to do structurally better:

Theorem (S. Cook, 1971; L. Levin, 1973)

The problem of deciding classical logic is co-NP-complete.

Łukasiewicz infinite-valued propositional logic



Jan Łukasiewicz, 1878-1956.

Łukasiewicz infinite-valued propositional logic



Jan Łukasiewicz, 1878-1956.

The formulæ of Łukasiewicz logic are the same as the ones in CL: FORM and $FORM_n$ are as before.

Axiom system for Łukasiewicz logic.

$$(A0) \perp \rightarrow \alpha \qquad \qquad Ex \ falso \ quod libet.$$

$$(A1) \ \alpha \rightarrow (\beta \rightarrow \alpha) \qquad \qquad A \ fortiori.$$

$$(A2) \ (\alpha \rightarrow \beta) \rightarrow ((\beta \rightarrow \gamma) \rightarrow (\alpha \rightarrow \gamma)) \qquad \text{Implication is transitive.}$$

$$(A3) \ (\alpha \lor \beta) \rightarrow (\beta \lor \alpha) \qquad \qquad \text{Disjunction is commutative.}$$

$$(A4) \ (\alpha \rightarrow \beta) \rightarrow (\neg \beta \rightarrow \neg \alpha) \qquad \qquad \text{Contraposition.}$$

$$(A5) \ \alpha \lor \neg \alpha \qquad \qquad \qquad \overline{Tertium \ non \ datur \ / Consequentia \ Mirabilis.}$$

Axiom system for Łukasiewicz logic.

(A0)
$$\perp \rightarrow \alpha$$
 Ex falso quodlibet.

(A1)
$$\alpha \rightarrow (\beta \rightarrow \alpha)$$
 A fortiori.

(A2)
$$(\alpha \to \beta) \to ((\beta \to \gamma) \to (\alpha \to \gamma))$$
 Implication is transitive.

(A3)
$$(\alpha \lor \beta) \to (\beta \lor \alpha)$$
 Disjunction is commutative.

$$\text{(A4)} \ (\alpha \to \beta) \ \to \ (\neg \beta \to \neg \alpha)$$
 Contraposition.

(A5)
$$\alpha \vee \neg \alpha$$
 Tertium non datur / Consequentia Mirabilis.

Deduction rule for Łukasiewicz logic.

(R1)
$$\frac{\alpha \qquad \alpha \rightarrow \beta}{\beta}$$

Modus ponens.

Lukasiewicz infinite-valued propositional logic

Łukasiewicz logic can be succinctly described as classical logic without the Aristotelian law of Tertium non datur, but with the Ex falso quodlibet law.

Łukasiewicz logic can be succinctly described as classical logic without the Aristotelian law of Tertium non datur, but with the Ex falso quodlibet law.

Such "succint descriptions" can be polysemous to a surprising extent indeed.

Łukasiewicz logic Intuitionistic logic can be succinctly described as classical logic without the Aristotelian law of Tertium non datur, but with the Ex falso quodlibet law.¹

Such "succint descriptions" can be polysemous to a surprising extent indeed.

¹Almost *verbatim* from J. Moschovakis, *Intuitionistic Logic*, The Stanford Encyclopedia of Philosophy, 2010, Edward N. Zalta (ed.).

Łukasiewicz logic Intuitionistic logic can be succinctly described as classical logic without the Aristotelian law of Tertium non datur, but with the Ex falso quodlibet law.¹

Such "succint descriptions" can be polysemous to a surprising extent indeed.

Moral: The import of removing an axiom from an axiom system depends on the axiom system itself. In particular, Hilbert-style systems are of little use to analyse the structural properties of logics in terms of a specific axiomatisation.

¹Almost *verbatim* from J. Moschovakis, *Intuitionistic Logic*, The Stanford Encyclopedia of Philosophy, 2010, Edward N. Zalta (ed.).

As for CL, we are using $\{\bot, \neg, \rightarrow\}$ as primitive connectives. We define $\lor, \land, \leftrightarrow, \ominus$ as before. It is customary to define more.

As for CL, we are using $\{\bot, \neg, \rightarrow\}$ as primitive connectives. We define $\lor, \land, \leftrightarrow, \ominus$ as before. It is customary to define more.

| Notation | Definition | Name | Idempotent |
|--------------------------------|--|-----------------------|------------|
| | _ | Falsum | _ |
| Т | $\neg \bot$ | Verum | _ |
| $\neg \alpha$ | _ | Negation | _ |
| $\alpha 	o \beta$ | _ | Implication | _ |
| $\alpha \vee \beta$ | $(\alpha \to \beta) \to \beta$ | (Lattice) Disjunction | Yes |
| α / β | $\neg(\neg\alpha\vee\neg\beta)$ | (Lattice) Conjunction | Yes |
| $\alpha \leftrightarrow \beta$ | $(\alpha \to \beta) \wedge (\beta \to \alpha)$ | Biconditional | _ |
| $\alpha \oplus \beta$ | eg lpha ightarrow eta | Strong disjunction | No |
| α ⊙ β | $\neg(\alpha \to \neg\beta)$ | Strong conjunction | No |
| $\alpha\ominus\beta$ | $\neg(\alpha \to \beta)$ | Co-implication | _ |

Table: Connectives in Łukasiewicz logic.

Lukasiewicz infinite-valued propositional logic

We now turn to semantics. Łukasiewicz logic has a many-valued semantics: specifically, pace Frege, we take $[0,1]\subseteq\mathbb{R}$ as a set of "truth values".

We now turn to semantics. Łukasiewicz logic has a many-valued semantics: specifically, *pace* Frege, we take $[0,1]\subseteq\mathbb{R}$ as a set of "truth values".

A possible world is an assignment

$$w \colon \text{Form} \to [0, 1]$$

We now turn to semantics. Łukasiewicz logic has a many-valued semantics: specifically, *pace* Frege, we take $[0,1] \subseteq \mathbb{R}$ as a set of "truth values".

A possible world is an assignment

$$w \colon \text{Form} \to [0, 1]$$

■
$$w(\bot) = 0$$
.

We now turn to semantics. Łukasiewicz logic has a many-valued semantics: specifically, *pace* Frege, we take $[0,1]\subseteq\mathbb{R}$ as a set of "truth values".

A possible world is an assignment

$$w \colon \text{Form} \to [0, 1]$$


- $w(\perp) = 0.$
- $w(\neg \alpha) = 1 w(\alpha).$

We now turn to semantics. Łukasiewicz logic has a many-valued semantics: specifically, *pace* Frege, we take $[0,1] \subseteq \mathbb{R}$ as a set of "truth values".

A possible world is an assignment


$$w \colon \text{Form} \to [0, 1]$$

- $w(\bot) = 0$.
- $w(\neg \alpha) = 1 w(\alpha).$



Truth-function of Łukasiewicz implication.

$$w(\alpha \to \beta) = \left\{ \begin{array}{ll} 1 & \text{if } w(\alpha) \leqslant w(\beta) \\ 1 - (w(\alpha) - w(\beta)) & \text{otherwise.} \end{array} \right.$$




Truth-function of Łukasiewicz implication.

$$w(\alpha \rightarrow \beta) = \min\{1, 1 - (w(\alpha) - w(\beta))\}\$$

| Notation | Formal semantics |
|--------------------------------|---|
| | $w(\perp) = 0$ |
| Т | $w(\top) = 1$ |
| $\neg \alpha$ | $w(\neg \alpha) = 1 - w(\alpha)$ |
| $\alpha 	o \beta$ | $w(\alpha \rightarrow \beta) = \min\{1, 1 - (w(\alpha) - w(\beta))\}$ |
| $\alpha \vee \beta$ | $w(\alpha \lor \beta) = \max\{w(\alpha), w(\beta)\}$ |
| α / β | $w(\alpha \wedge \beta) = \min\{w(\alpha), w(\beta)\}$ |
| $\alpha \leftrightarrow \beta$ | $w(\alpha \leftrightarrow \beta) = 1 - w(\alpha) - w(\beta) $ |
| $\alpha \oplus \beta$ | $w(\alpha \oplus \beta) = \min\{1, w(\alpha) + w(\beta)\}$ |
| $\alpha \odot \beta$ | $w(\alpha \odot \beta) = \max\{0, w(\alpha) + w(\beta) - 1\}$ |
| $\alpha\ominus\beta$ | $w(\alpha \ominus \beta) = \max\{0, w(\alpha) - w(\beta)\}$ |

Table: Formal semantics of connectives in Łukasiewicz logic.



 $\begin{tabular}{ll} Truth-function of Lukasiewicz "strong conjunction" \odot. \\ & (\textit{Note: Non-idempotent operation.}) \end{tabular}$

$$w(\alpha \odot \beta) = \max\{0, w(\alpha) + w(\beta) - 1\}$$

As in classical logic, a tautology is a formula α that receives value 1 in every possible world, i.e. for every assignment $w \colon \text{FORM} \to [0,1]$ it so happens that $w(\alpha) = 1$. So 1 is the only designated value.

As in classical logic, a tautology is a formula α that receives value 1 in every possible world, i.e. for every assignment $w \colon \mathrm{FORM} \to [0,1]$ it so happens that $w(\alpha) = 1$. So 1 is the only designated value.

Similarly, α is provable if it can be derived (in the previous sense) from the axioms of Łukasiewicz logic using *modus ponens*.

As in classical logic, a tautology is a formula α that receives value 1 in every possible world, i.e. for every assignment $w \colon \mathrm{FORM} \to [0,1]$ it so happens that $w(\alpha) = 1$. So 1 is the only designated value.

Similarly, α is provable if it can be derived (in the previous sense) from the axioms of Łukasiewicz logic using *modus ponens*.

Notation.

As in classical logic, a tautology is a formula α that receives value 1 in every possible world, i.e. for every assignment $w : \text{Form} \to [0, 1]$ it so happens that $w(\alpha) = 1$. So 1 is the only designated value.

Similarly, α is provable if it can be derived (in the previous sense) from the axioms of Łukasiewicz logic using modus ponens.

Notation.

- $\vdash_{i} \alpha$
- $\models_i \alpha$
- T_{HM}^{L}
- TAUTL

- α is provable.
- α is a tautology.
- Set of provable formulæ.
 - Set of tautologies.

Similarly for THM_n^L and $TAUT_n^L$.

Lukasiewicz infinite-valued propositional logic

Easy to show:

Easy to show:

Validity or Soundness Theorem for Ł

For any $\alpha \in \mathrm{FORM}$,

$$\vdash_{\mathsf{L}} \alpha \text{ implies } \vDash_{\mathsf{L}} \alpha.$$

That is,

$$\mathrm{Thm}^{\mathbf{L}} \subseteq \mathrm{Taut}^{\mathbf{L}}$$
.

(Similarly, $THM_n^{\mathsf{L}} \subseteq TAUT_n^{\mathsf{L}}$).

Lukasiewicz infinite-valued propositional logic

How do Łukasiewicz theorems and tautologies relate to the classical ones?

Clearly: $THM^{\mathsf{L}} \subseteq THM$, and similarly $THM_n^{\mathsf{L}} \subseteq THM_n$.

Clearly: $THM^{\mathsf{L}} \subseteq THM$, and similarly $THM_n^{\mathsf{L}} \subseteq THM_n$.

Clearly: $TAUT^{\ell} \subseteq TAUT$, and similarly $TAUT^{\ell}_n \subseteq THM_n$.

Clearly: $THM^{\mathsf{L}} \subseteq THM$, and similarly $THM_n^{\mathsf{L}} \subseteq THM_n$.

Clearly: $\mathrm{TAUT}^{\mathsf{L}} \subseteq \mathrm{TAUT}$, and similarly $\mathrm{TAUT}_n^{\mathsf{L}} \subseteq \mathrm{THM}_n$.

The latter (semantic) inclusion is proper: there is a classical tautology that is not a Łukasiewicz tautology, namely, the law of excluded middle.

Clearly: $THM^{\mathsf{L}} \subseteq THM$, and similarly $THM_n^{\mathsf{L}} \subseteq THM_n$.

Clearly: $\mathrm{TAUT}^{\mathsf{L}} \subseteq \mathrm{TAUT}$, and similarly $\mathrm{TAUT}_n^{\mathsf{L}} \subseteq \mathrm{THM}_n$.

The latter (semantic) inclusion is proper: there is a classical tautology that is not a Łukasiewicz tautology, namely, the law of excluded middle.

$$\not\models_{\mathsf{L}} p \lor \lnot p$$

(*Proof.* Take $w(p) = \frac{1}{2}$.)

Clearly: $THM^{\mathsf{L}} \subseteq THM$, and similarly $THM_n^{\mathsf{L}} \subseteq THM_n$.

Clearly: $\mathrm{TAUT}^{\mathsf{L}} \subseteq \mathrm{TAUT}$, and similarly $\mathrm{TAUT}_n^{\mathsf{L}} \subseteq \mathrm{THM}_n$.

The latter (semantic) inclusion is proper: there is a classical tautology that is not a Łukasiewicz tautology, namely, the law of excluded middle.

$$\not\models_{\mathsf{L}} p \lor \lnot p$$

(*Proof.* Take $w(p) = \frac{1}{2}$.) In fact, we have seen "by construction" that

$$\mathbf{L} + (\alpha \vee \neg \alpha) = CL$$
.

Clearly: $THM^{\mathsf{L}} \subseteq THM$, and similarly $THM_n^{\mathsf{L}} \subseteq THM_n$.

Clearly: $\mathrm{TAUT}^{\mathsf{L}} \subseteq \mathrm{TAUT}$, and similarly $\mathrm{TAUT}_n^{\mathsf{L}} \subseteq \mathrm{THM}_n$.

The latter (semantic) inclusion is proper: there is a classical tautology that is not a Łukasiewicz tautology, namely, the law of excluded middle.

$$\not\models_{\mathsf{L}} p \lor \lnot p$$

(*Proof.* Take $w(p) = \frac{1}{2}$.) In fact, we have seen "by construction" that

$$L + (\alpha \vee \neg \alpha) = CL$$
.

The (syntactic) inclusion $\mathrm{THM}^{\mathsf{L}} \subseteq \mathrm{THM}$ also is proper. The full relationship between tautologies and theorems follows from the next result, a substantial piece of mathematics.

Completeness Theorem for Ł (A. Rose and J. B. Rosser, 1958)

For any $\alpha \in \mathrm{FORM}$,

$$\models_{\mathsf{L}} \alpha$$
 implies $\vdash_{\mathsf{L}} \alpha$.

That is,

$$\text{TAUT}^{\mathsf{L}} \subseteq \text{THM}^{\mathsf{L}}$$
.

Completeness Theorem for Ł (A. Rose and J. B. Rosser, 1958)

For any $\alpha \in \mathrm{FORM}$,

$$\models_{\mathsf{L}} \alpha$$
 implies $\vdash_{\mathsf{L}} \alpha$.

That is,

$$TAUT^{L} \subseteq THM^{L}$$
.

(Similarly, TAUT_n \subseteq THM_n.).

Lukasiewicz infinite-valued propositional logic

In summary:

In summary:

Soundness and Completeness Theorem for Ł

For any $\alpha \in \mathrm{FORM}$,

$$\models_{\mathsf{L}} \alpha$$
 if, and only if, $\vdash_{\mathsf{L}} \alpha$.

That is,

$$\mathrm{TAUT}^{\boldsymbol{\mathsf{L}}} = \mathrm{THM}^{\boldsymbol{\mathsf{L}}}$$
 .

(Similarly,
$$TAUT_n^{\mathsf{L}} = THM_n^{\mathsf{L}}$$
).

² Added in proof. We did not have time to sketch the proof of this result.

Corollary (D. Mundici, 1987)

There is a an algorithm (for definiteness, a Turing machine) that, on input $\alpha \in \mathrm{FORM}$, outputs YES if $\alpha \in \mathrm{THM}^L$, and NO otherwise.

² Added in proof. We did not have time to sketch the proof of this result.

Corollary (D. Mundici, 1987)

There is a an algorithm (for definiteness, a Turing machine) that, on input $\alpha \in \mathrm{FORM}$, outputs YES if $\alpha \in \mathrm{THM}^L$, and NO otherwise.

But even with completeness, a brute-force check of all [0,1]-valued assignments to the n variables in α is impossible. Something cleverer is needed (we will perhaps hint at a proof later).²

² Added in proof. We did not have time to sketch the proof of this result.

Corollary (D. Mundici, 1987)

There is a an algorithm (for definiteness, a Turing machine) that, on input $\alpha \in \mathrm{FORM}$, outputs YES if $\alpha \in \mathrm{THM}^L$, and No otherwise.

But even with completeness, a brute-force check of all [0,1]-valued assignments to the n variables in α is impossible. Something cleverer is needed (we will perhaps hint at a proof later).²

And yet, structurally, the situation is no worse than in the classical case:

Theorem (D. Mundici, 1987)

The problem of deciding Łukasiewicz logic is co-NP-complete.

² Added in proof. We did not have time to sketch the proof of this result.

Lukasiewicz infinite-valued propositional logic

Thank you for your attention.